

p21 (phospho-T145) Antibody

Basic information:

Catalog No.:UPA02841Source:RabbitSize:50ul/100ulClonality:Polyclonal

Concentration: 1mg/ml Isotype: IgG

Purification: The antibody was purified by immunogen affinity chromatography.

Useful Information:

WB: 1:500 - 1:1000

Applications: IHC: 1:100-1:200

IF/IC: 1:100-1:500

Reactivity: Human, Bovine, Dog, Sheep

Specificity: Recognizes endogenous levels of p21 (pT145) protein.

KLH-conjugated synthetic peptide encompassing a sequence within the **Immunogen:**

C-term region of human p21 (pT145). The exact sequence is proprietary.

Description: Rabbit polyclonal antibody to p21 (pT145)

Uniprot: P38936(Human)
BiowMW: Refer to Figures


Buffer: Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30%

glycerol, and 0.01% sodium azide.

Storage: Store at 4°C short term and -20°C long term. Avoid freeze-thaw cycles.

Note: For research use only, not for use in diagnostic procedure.

Data:




Western blot analysis of p21 (phospho-T145) expression in HT1080 EGF-treated (A), HeLa EGF-treated (B) whole cell lysates.


E-mail: sales@universalbiol.com

Tel: 0550-3121009









Immunohistochemical analysis of p21 (phospho-T145) staining in human prostate cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Immunofluorescent analysis of p21 (phospho-T145) staining in HeLa cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a hidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark.