

p67 Antibody


Basic information:

Catalog No.:	UPA03842	Source:	Rabbit
Size:	50ul/100ul	Clonality:	Polyclonal
Concentration:	1mg/ml	Isotype:	IgG
Purification:	The antibody was purified by immunogen affinity chromatography.		


Useful Information:

Applications:	WB (1:500 - 1:1000), IHC (1:100 - 1:200), IF/ICC (1:100 - 1:500)
Reactivity:	Human, Mouse, Rat, Bovine
Specificity:	Recognizes endogenous levels of p67 protein.
Immunogen:	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human p67. The exact sequence is proprietary.
Description:	Rabbit polyclonal antibody to p67
Uniprot:	P61764
BiowMW:	Refer to Figures
Buffer:	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% Thimerosal.
Storage:	Shipped at 4°C. Upon delivery aliquot and store at -20°C for one year. Avoid freeze/thaw cycles.
Note:	For research use only, not for use in diagnostic procedure.


Data:



Western blot analysis of p67 expression in U87MG (A), C6 (B), SHSY5Y (C), K562 (D), Raw264.7 (E), mouse brain (F), rat brain (G) whole cell lysates. (Predicted band size: 67 kD; Observed band size: 70 kD)



Immunohistochemical analysis of p67 staining in human brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Immunofluorescent analysis of p67 staining in Jurkat cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).